Decreasing Narcotic Prescribing One Urban ED's Experience

MNACEP Leadership Conference January 28, 2019


Chris Palmer, MD, FACEP
North Memorial ED Medical Director
EPPA Physician

Disclosures: None

Special Thanks:

Megan Blazevic Cameron Berg MD NM/MG ED Providers

Opioid Epidemic










Opioid Overdoses









Opioid Impact

In 2016, there were 395 opioid overdose deaths. For every 1 death there were...

Death



† 1

Hospital admissions



†† 2

Emergency department visits





EMS responses














It's Illegal Drugs Right?? - WRONG

Rates of Prescription Painkiller Sales, Deaths, and Substance Abuse Treatment Admissions (1999-2010)






Prescription Painkiller Overdoses in the US; CDC Vital Signs November 201









mg of prescription opioids sold




nillions of milligrams





A Prescribing Problem

Number of Opioid Prescriptions Dispensed by U.S. Retail Pharmacies, Years 1991-2011










When Does the Risk Start?



Days' supply of first opioid prescription


Days' supply of first opioid prescription







When Does the Risk Start?










Which Specialty

Top 10 specialties by opioid prescription volume (2016)









New England Journal of Medicine Study

- Opioid Prescribing Patterns of Emergency Physicians and Risk of Long-Term Use
 - Michael L. Barnett, M.D., Andrew R. Olenski, B.S., and Anupam B. Jena, M.D., Ph.D.
- NEJM: Feb 2017
- Basic conclusion: Introduction to opioids matters







Public Education

- Public health dedication (National and Local Media Attention)
- Mass media focus
- Population more aware
- Narcan kits
- Legislative funding







Institute for Clinical Systems Improvement (ICSI)

State collaboration to tackle this issue

- Agreement that 100MEQ is a threshold that increases risk of long term addiction
- NM/MG ED providers were previously focused on this work
- NM/MG ED providers added this to Quality Improvement work in 2018








Process

- Developed measurement tools for individuals
- Published blinded information quarterly
- Discussed at team meetings and in performance evaluations
- Reviewed relevant best practice and published evidence
- Reviewed progress annually

























ICSI (Recommendations late 2017)

- ICSI identified 100MME as concerning threshold
- Baseline at NM/MG: 1/1/2016 through 12/4/2017 3.8% of ED discharges had opioid prescription exceeding 100 MME
- Impact: Reducing unnecessary or inappropriate prescriptions exceeding 100 MME will confer improved safety to our customers.
- Target: ≤3.0% (21% relative reduction)
- Result: NM/MG providers reduced prescriptions
 >100 MME to 0.3%
























Results: 2018 changes compared to 2015 baseline

- 2.146 million fewer MME dispensed
 - 429,210 less hydrocodone (5mg) or 286,140 less oxycodone (5 mg)
- Reduction from 26.1 MME/pt seen to 6.33 MME/pt seen
- Average prescription from 142.87 to 92.06 MME
- 17,080 fewer opiate prescriptions
- 10,920 fewer opiate scripts exceeding 100 MME







Results: 2018 changes compared to 2015 baseline

- 15,384 fewer opiate prescriptions
- 10,450 fewer opiate scripts exceeding 100 MME
- 1,981 fewer opiate scripts exceeding 200 MME

It comes down to individuals

- Unique impact
- Decision to change





